

- [0124] 43. Elnakat H, Ratnam M: Distribution, functionality and gene regulation of folate receptor isoforms: implications in targeted therapy, *Advanced drug delivery reviews* 2004, 56:1067-1084
- [0125] 44. Parker N, Turk M J, Westrick E, Lewis J D, Low P S, Leamon C P: Folate receptor expression in carcinomas and normal tissues determined by a quantitative radioligand binding assay, *Analytical biochemistry* 2005, 338:284-293
- [0126] 45. Weitman S D, Lark R H, Coney L R, Fort D W, Frasca V, Zurawski V R, Jr., Kamen B A: Distribution of the folate receptor GP38 in normal and malignant cell lines and tissues, *Cancer research* 1992, 52:3396-3401
- [0127] 46. Kane M A, Elwood P C, Portillo R M, Antony A C, Najfeld V, Finley A, Waxman S, Kolhouse J F: Influence on immunoreactive folate-binding proteins of extracellular folate concentration in cultured human cells, *The Journal of clinical investigation* 1988, 81:1398-1406
- [0128] 47. Cho B K, Roy E J, Patrick T A, Kranz D M: Single-chain Fv/folate conjugates mediate efficient lysis of folate-receptor-positive tumor cells, *Bioconjugate chemistry* 1997, 8:338-346
- [0129] 48. Speyer C L, Gao H, Rancilio N J, Neff T A, Huffnagle G B, Sarma J V, Ward P A: Novel chemokine responsiveness and mobilization of neutrophils during sepsis, *The American journal of pathology* 2004, 165: 2187-2196
- [0130] 49. Ethier S P, Mahacek M L, Gullick W J, Frank T S, Weber B L: Differential isolation of normal luminal mammary epithelial cells and breast cancer cells from primary and metastatic sites using selective media, *Cancer research* 1993, 53:627-635
- [0131] 50. Tran T, Shatnawi A, Zheng X, Kelley K M, Ratnam M: Enhancement of folate receptor alpha expression in tumor cells through the glucocorticoid receptor: a promising means to improved tumor detection and targeting, *Cancer research* 2005, 65:4431-4441
- [0132] 51. (king-Konert C, Vogt S, Radsak M, Wagner C, Hansch G M, Andrassy K: Polymorphonuclear neutrophils in Wegener's granulomatosis acquire characteristics of antigen presenting cells, *Kidney Int* 2001, 60:2247-2262
- [0133] 52. van Egmond M, van Vuuren A J, Morton H C, van Spruiel A B, Shen L, Hofhuis F M, Saito T, Mayadas T N, Verbeek J S, van de Winkel J G: Human immunoglobulin A receptor (FcalphaRI, CD89) function in transgenic mice requires both FcR gamma chain and CR3 (CD11b/CD18), *Blood* 1999, 93:4387-4394
- [0134] 53. Weisbart R H, Kacena A, Schuh A, Golde D W: GM-CSF induces human neutrophil IgA-mediated phagocytosis by an IgA Fc receptor activation mechanism, *Nature* 1988, 332:647-648
- [0135] 54. Morton H C, van Egmond M, van de Winkel J G: Structure and function of human IgA Fc receptors (Fc alpha R), *Critical reviews in immunology* 1996, 16:423-440
- [0136] 55. Monteiro R C, Van De Winkel J G: IgA Fc receptors, *Annu Rev Immunol* 2003, 21:177-204
- [0137] 56. Reljic R: In search of the elusive mouse macrophage Fc-alpha receptor, *Immunol Lett* 2006, 107:80-81
- [0138] 57. Prabakaran M, Grailer J J, Pilla S, Steeber D A, Gong S: Gold nanoparticles with a monolayer of doxorubicin-conjugated amphiphilic block copolymer for tumor-targeted drug delivery, *Biomaterials* 2009, 30:6065-6075
- [0139] 58. Prabakaran M, Grailer J J, Pilla S, Steeber D A, Gong S: Folate-conjugated amphiphilic hyperbranched block copolymers based on Boltorn H40, poly(L-lactide) and poly(ethylene glycol) for tumor-targeted drug delivery, *Biomaterials* 2009, 30:3009-3019
- [0140] 59. Prabakaran M, Grailer J J, Steeber D A, Gong S: Thermosensitive micelles based on folate-conjugated poly(N-vinylcaprolactam)-block-poly(ethylene glycol) for tumor-targeted drug delivery, *Macromolecular bioscience* 2009, 9:744-753
- [0141] 60. Li H, Lu Y, Piao L, Wu J, Yang X, Kondadasula S V, Carson W E, Lee R J: Folate-immunoglobulin G as an anticancer therapeutic antibody, *Bioconjugate chemistry* 2010, 21:961-968
- [0142] 61. Speyer C L, Smith J S, Banda M, DeVries J A, Mekani T, Gorski D H: Metabotropic glutamate receptor-1: a potential therapeutic target for the treatment of breast cancer, *Breast cancer research and treatment* 2012, 132: 565-573
- What is claimed is:
1. A conjugate comprising:  
an Fc portion of an IgA antibody, wherein said Fc portion comprises a CH2 domain and a CH3 domain but does not comprise a hinge region; and  
folate or a folate analog.
  2. The conjugate of claim 1, further comprising a linker.
  3. The conjugate of claim 2, wherein the linker is a peptide linker.
  4. The conjugate of claim 2, wherein the linker is a non-peptide linker.
  5. The conjugate of claim 1, wherein the Fc portion is a variant Fc.
  6. The conjugate of claim 1, wherein the Fc portion is a variant Fc and the conjugate comprises folate.
  7. The conjugate of claim 1, wherein the Fc portion is a variant Fc and the conjugate comprises a folate analog.
  8. The conjugate of claim 1, wherein the Fc fragment consists of a CH2/CH3 region from an IgA antibody.
  9. The conjugate of claim 1, formulated for intravenous administration.
  10. A method of triggering neutrophil (PMN)-mediated apoptosis of folate receptor alpha (FRA)-expressing cancer cells in a subject in need thereof, comprising: administering to the subject a therapeutically effective amount of the conjugate of claim 1, thereby triggering neutrophil (PMN)-mediated apoptosis of folate receptor alpha (FRA)-expressing cancer cells in the subject.
  11. The method of claim 10, wherein the cancer is breast cancer, ovarian cancer, or lung cancer.
  12. The method of claim 10, wherein the cancer is triple negative breast cancer.
  13. The method of claim 10, wherein the cancer includes cancer cells that lack receptors for one or more of estrogen, progesterone and epithelial growth factor.
  14. The method of claim 10, wherein administration is by intravenous administration.
  15. A method of recruiting and activating neutrophils to destroy a folate receptor alpha (FRA)-expressing cancer cell in a subject, comprising contacting the cancer cell with a conjugate comprising: